



Marshall, A. D., Cunningham, E. B., Nielsen, S., Aghemo, A., Alho, H., Backmund, M., Bruggmann, P., Dalgard, O., Seguin-Devaux, C., Flisiak, R., Foster, G. R., Gheorghe, L., Goldberg, D., Goulis, I., Hickman, M., Hoffmann, P., Jancoriené, L., Jarcuska, P., Kåberg, M. (2018). Restrictions for reimbursement of interferon-free direct-acting antiviral drugs for HCV infection in Europe. *Lancet Gastroenterology and Hepatology*, 3(2), 125–133. [https://doi.org/10.1016/S2468-1253\(17\)30284-4](https://doi.org/10.1016/S2468-1253(17)30284-4)

Peer reviewed version

License (if available):  
CC BY-NC-ND

Link to published version (if available):  
[10.1016/S2468-1253\(17\)30284-4](https://doi.org/10.1016/S2468-1253(17)30284-4)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via The Lancet Group at <https://www.sciencedirect.com/science/article/pii/S2468125317302844> . Please refer to any applicable terms of use of the publisher.

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

# **Restrictions for reimbursement of interferon-free direct-acting antiviral therapies for HCV infection in Europe**

Alison D. Marshall (MA)<sup>1</sup>, Evan B. Cunningham (BSc Hons)<sup>1</sup>, Stine Nielsen (MSc)<sup>2</sup>, Alessio Aghemo ( PhD)<sup>3,4</sup>, Hannu Alho (Prof, PhD)<sup>5</sup>, Markus Backmund ( PhD)<sup>6</sup>, Philip Bruggmann (MD)<sup>7</sup>, Olav Dalgard (PhD)<sup>8</sup>, Carole Seguin-Devaux (PhD)<sup>9</sup>, Robert Flisiak ( PhD)<sup>10</sup>, Graham R. Foster ( PhD)<sup>11</sup>, Liana Gheorghe ( PhD)<sup>12</sup>, David Goldberg (MD)<sup>13</sup>, Ioannis Goulis (MD)<sup>14</sup>, Matthew Hickman (Prof, PhD)<sup>15</sup>, Patrick Hoffmann (PhD)<sup>16</sup>, Ligita Jancorienė (MD, PhD)<sup>17</sup>, Peter Jarcuska ( PhD)<sup>18</sup>, Martin Kåberg (MD)<sup>19</sup>, Leondios G. Kostrikis (PhD)<sup>20</sup>, Mihály Makara (MD)<sup>21</sup>, Matti Maimets ( PhD)<sup>22</sup>, Rui Tato Marinho ( PhD)<sup>23</sup>, Mojca Matičič (Prof, , PhD)<sup>24</sup>, Suzanne Norris ( PhD)<sup>25</sup>, Sigurður Ólafsson (MD)<sup>26</sup>, Anne Øvrehus (MD)<sup>27</sup>, Jean-Michel Pawlotsky ( PhD)<sup>28</sup>, James Pocock (MD)<sup>29</sup>, Geert Robaeys ( PhD)<sup>30,31,32</sup>, Carlos Roncero ( PhD)<sup>33</sup>, Marieta Simonova ( PhD)<sup>34</sup>, Jan Sperl ( PhD)<sup>35</sup>, Michele Tait (BSc)<sup>25</sup>, Ieva Tolmane ( PhD)<sup>36,37</sup>, Stefan Tomaselli (MSocEcSc)<sup>38</sup>, Marc van der Valk ( PhD)<sup>39</sup>, Adriana Vince (Prof, , PhD)<sup>40</sup>, Gregory J. Dore (Prof, PhD)<sup>1</sup>, Jeffrey V. Lazarus (PhD)<sup>41,42\*</sup>, and Jason Grebely (PhD)<sup>1\*</sup> on behalf of the International Network on Hepatitis in Substance Users (INHSU) Network

\*contributed equally

<sup>1</sup>The Kirby Institute, UNSW Sydney, NSW Australia <sup>2</sup>Freelance epidemiologist <sup>3</sup>Humanitas University Department of Biomedical Sciences, Milan, Italy <sup>4</sup>Humanitas Clinical and Research Center, Milan, Italy <sup>5</sup>Abdominal Center, Helsinki University Hospital and Helsinki University, Finland <sup>6</sup>Ludwig-Maximilians-University, Munich, Germany <sup>7</sup>Arud Centres for Addiction Medicine, Zürich, Switzerland <sup>8</sup>Department of Infectious Diseases, Akershus University Hospital and Faculty of Medicine, University of Oslo, Norway <sup>9</sup>Department of Infection and Immunity, Luxembourg Institute of Health, Luxembourg <sup>10</sup>Department of

Infectious Diseases and Hepatology, Medical University of Bialystok, Poland <sup>11</sup>Queen Mary, University of London, UK <sup>12</sup>Gastroenterology and Hepatology, Fundeni Clinical Institute, Bucharest, Romania <sup>13</sup>Health Protection Scotland (HPS), Glasgow, UK <sup>14</sup>Department of Internal Medicine, Thessaloniki, Greece <sup>15</sup>Social Medicine, University of Bristol, UK <sup>16</sup>Ministry of Health Luxembourg, Luxembourg City, Luxembourg <sup>17</sup>Centre of Infectious Diseases, Vilnius University Hospital Santaros Klinikos, Vilnius University, Vilnius, Lithuania <sup>18</sup>1st Department of Internal Medicine, University Hospital and University of Pavol Jozef Safarik, Faculty of Medicine, Kosice, Slovakia <sup>19</sup>Department of Infectious Diseases, Karolinska University Hospital, Stockholm, Sweden <sup>20</sup>University of Cyprus, Cyprus <sup>21</sup>St István and St László Hospital, Hepatology Center, Budapest, Hungary <sup>22</sup>Department of Internal Medicine, University of Tartu, Estonia <sup>23</sup>Department of Gastroenterology and Hepatology, Hospital Santa Maria, Medical School Lisbon, University of Lisbon, Portugal <sup>24</sup>Clinic for Infectious Diseases and Febrile Illnesses, University Medical Centre, Ljubljana, Slovenia <sup>25</sup>National Hepatitis C Treatment Programme, Health Service Executive, Dr Steevens' Hospital, Dublin, Ireland <sup>26</sup>Division of Gastroenterology, Department of Medicine, Landspítali University Hospital, Reykjavik, Iceland <sup>27</sup>Department of Infectious Diseases, Odense University Hospital, University of Southern Denmark, Odense, Denmark <sup>28</sup>Hopital Henri Mondor, Université Paris Est, Creteil, France <sup>29</sup>Gastroenterology Department, Mater Dei Hospital, Msida, Malta <sup>30</sup>Department of Gastroenterology and Hepatology, Ziekenhuis Oost Limburg, Genk, Belgium <sup>31</sup>Department of Medicine and Life Sciences, UHasselt, Hasselt, Belgium <sup>32</sup>Department of Hepatology, UZLeuven, Leuven, Belgium <sup>33</sup>Addiction and Dual Diagnosis Unit, Psychiatry Service, Hospital Universitario Vall d'Hebron. Psychiatric Department. Universidad Autónoma de Barcelona, Barcelona, Spain <sup>34</sup>Department of Gastroenterology, HPB Surgery and Transplantology, Military Medical Academy, Sofia, Bulgaria <sup>35</sup>Department of Hepatogastroenterology, Institute for Clinical and

Experimental Medicine, Prague, Czech Republic <sup>36</sup>Department of Hepatology, Infectology  
Center of Latvia, Riga East University Hospital, Riga, Latvia <sup>37</sup>University of Latvia, Faculty  
of Medicine <sup>38</sup>Office of Public Health, Vaduz, Liechtenstein <sup>39</sup>Department of Infectious  
Diseases, Academic Medical Center, Amsterdam, Netherlands <sup>40</sup>University Hospital of  
Infectious Diseases, University of Zagreb, School of Medicine, Zagreb, Croatia <sup>41</sup>Barcelona  
Institute of Global Health (ISGlobal), Hospital Clínic, Barcelona, Spain <sup>42</sup>CHIP,  
Rigshospitalet, University of Copenhagen, Denmark

**Corresponding author**

Alison Marshall

Viral Hepatitis Clinical Research Program

The Kirby Institute

UNSW Sydney, Sydney, Australia

Phone: +61-2-9385 9960 Fax: +61-2-938 5 0876

Email: [amarshall@kirby.unsw.edu.au](mailto:amarshall@kirby.unsw.edu.au)

Abstract word count: 150

Manuscript word count: 3,737

Figures and Tables: 5

References: 61

## Abstract

All-oral direct-acting antiviral (DAA) HCV therapies with cure responses of >90% is a major clinical advancement. However, the high DAA list price has led many governments to restrict their reimbursement. Study aims were to review the availability of, and national criteria for, interferon-free DAA therapy reimbursement among EU/EEA countries and Switzerland. Reimbursement documentation was reviewed from November 18, 2016 to August 1, 2017. Primary outcomes were fibrosis stage, drug or alcohol use, prescriber type, and HIV co-infection restrictions. Among European countries/jurisdictions (n=35), the most commonly reimbursed DAA was ombitasvir/paritaprevir/ritonavir  $\pm$  dasabuvir  $\pm$  ribavirin (94%, n=33), nearly half of countries/jurisdictions (49%, n=17) required  $\geq$ F2, 83% (n=29) had no listed drug or alcohol use restrictions, 94% (n=33) required a specialist prescriber, and 97% (n=34) had no additional restrictions for HIV-HCV co-infected persons. Findings have implications for meeting WHO targets with evidence of some countries not following the 2016 EASL HCV treatment guidelines. **Funding:** The Kirby Institute is funded by the Australian Government Department of Health. The views expressed in this publication do not necessarily represent the position of the Australian Government.

**Key words:** hepatitis C virus, hepatitis C treatment, direct-acting antiviral, treatment restrictions, reimbursement, Europe, liver fibrosis, alcohol use, PWID, HIV-HCV co-infection

## INTRODUCTION

Over 71 million (63 – 79) people are infected with chronic hepatitis C virus (HCV) infection globally with ~704,000 (652,100 – 769,600) HCV-related deaths each year.<sup>1-3</sup> Compared to peginterferon-based therapies, simple, tolerable, all-oral HCV direct-acting antiviral (DAA) therapies achieve viral cure in  $\geq 90\%$  of patients, resulting in one of the greatest clinical advances in recent decades.<sup>4</sup> Broad uptake of DAA therapies has the potential to substantially reduce the global HCV disease burden, in particular, HCV-related mortality and morbidity.<sup>5-7</sup> However, the high list price of DAA therapies in some countries has led national governments<sup>8,9</sup> to restrict patient reimbursement based on liver fibrosis severity, drug and alcohol use, and prescriber type.

A 2014 study of Medicaid reimbursement criteria for sofosbuvir in the United States (US) showed some variability by state jurisdiction.<sup>8</sup> The majority of states (88%) had drug and/or alcohol use restrictions, most (74%) required evidence of advanced fibrosis ( $\geq F3$ ), over-half (66%) had prescriber limitations, and one-quarter (25%) required HIV-HCV co-infected persons to demonstrate suppressed HIV RNA levels or be recipients of antiretroviral therapy.<sup>8</sup> An updated reimbursement study by Ooka et al. (2017) demonstrated that, overall, restrictions had lessened over a two year period (2014 - 2016).<sup>10</sup> For example, 22 states required evidence of  $\geq F3$  in 2016 as opposed to 31 states in 2014. This updated study highlights that several criteria continue to be incongruent with clinical recommendations which state that all persons willing to be treated (and who have no treatment contraindications) should be considered for HCV therapy.<sup>11-13</sup>

In Europe, ~3.2 million (2.1 – 3.8) persons are estimated to have chronic HCV infection.<sup>14</sup> The primary HCV transmission route in the majority of European countries is injection drug

use.<sup>15-17</sup> Although HCV incidence is decreasing overall in Europe, rates of liver-related deaths, decompensated cirrhosis, and hepatocellular carcinoma are all projected to increase 20-30% by 2030 if broad access to DAA therapy is not achieved.<sup>14</sup> This, in the context of 2030 HCV global targets set by the World Health Organization (WHO) which propose: a 80% reduction in HCV incidence, 90% of persons with HCV diagnosed, 80% of persons with HCV treated, and a 65% reduction in HCV-related mortality.<sup>18</sup> To meet WHO targets, it will be critical that countries incorporate strategies that optimise uptake of DAA therapy, including minimising DAA reimbursement restrictions.

The aims of the study were to review the availability of interferon-free DAA therapy among European Union and European Economic Area (EU/EEA) countries and Switzerland; and to review national criteria for DAA therapy reimbursement among EU/EEA countries and Switzerland.



## METHODOLOGY

### *Data collection*

This study reviewed the availability of reimbursed DAA therapy and national criteria for DAA therapy reimbursement criteria among EU/EEA countries (including England, Northern Ireland, Scotland and Wales as separate jurisdictions) and Switzerland. Reimbursement criteria were reviewed for the following DAA regimens: sofosbuvir + ribavirin, sofosbuvir/ledipasvir  $\pm$  ribavirin, sofosbuvir/velpatasvir  $\pm$  ribavirin, ombitasvir/paritaprevir/ritonavir  $\pm$  dasabuvir  $\pm$  ribavirin, elbasvir-grazoprevir  $\pm$  ribavirin, and sofosbuvir + daclatasvir  $\pm$  ribavirin.

From November 18, 2016 to August 1, 2017, most data were extracted from publicly available documentation from government websites and online drug formularies with the assistance of study authors [see appendix p.1]. Supporting documentation, e.g. national HCV guidelines, was also collected when available [see appendix p.3]. If challenges arose in recovering information, one of three study authors contacted the study author from the country of interest to enable access. Given that most documentation was in a language other than English, study authors were fundamental to identifying documents of relevance. Independent document searches were conducted by three study authors to further corroborate search findings. Study authors of each country were also contacted to provide clarification if there were inconsistencies between documents and/or additional documentation was needed. In the event that documentation for a country could not be located, this finding was categorised as ‘Not available’.

Information was reviewed to determine the availability of reimbursed HCV DAAs in Europe (categories were: Yes; No). In the event that an individual could retrieve access to a specific

therapy (e.g. sofosbuvir + daclatasvir  $\pm$  ribavirin) but this therapy was not reimbursed, this was categorised as 'No'. Based on prior HCV DAA reimbursement studies,<sup>8,9</sup> primary outcomes were criteria restrictions based on: 1) minimum fibrosis stage [METAVIR or equivalent; Meta-Analysis of Histologic Data in Viral Hepatitis]; 2) drug and/or alcohol use; 3) prescriber type; and 4) HIV co-infection. If information for a specific outcome was not reported with the remaining outcomes, this outcome was categorised as 'None listed'.

Regarding primary outcomes, fibrosis stage was categorised as: No restriction; F2; F3, F4; Other; None listed; Not available. Fibrosis stage could be assessed through liver biopsy, transient elastography, biomedical markers, or a combination. It was anticipated that in many countries, patients with extrahepatic manifestations (e.g. cryoglobulinemia) would have had different criteria than asymptomatic patients (e.g. different fibrosis stage restrictions). Given that the primary interest of this study was to assess criteria experienced by the majority of patients, the minimum fibrosis stage for asymptomatic patients was recorded. Drug or alcohol use was categorised as: Prioritised; No restrictions; Additional restrictions; None listed; Not available. 'Prioritised' meant that persons with drug or alcohol use dependencies had fewer restrictions for DAA therapy access than persons without drug or alcohol use dependencies. 'No restrictions' meant that persons with drug or alcohol use dependencies could receive reimbursed DAA therapies. For instance, a person who engaged in active injection drug use (or a history of injection use) could be offered DAA therapy. 'Additional restrictions' meant that persons with drug or alcohol use dependencies needed to fulfil further criteria prior to being eligible for DAA therapy. Based on prior evidence,<sup>8, 10</sup> study authors anticipated that one example of a drug or alcohol use restriction that may be in existence in European countries would be a mandatory period of abstinence from substance use prior to therapy (e.g. 6 month period of abstinence from drug use). In addition to this restriction, study authors made note of any other drug or alcohol use restriction that was listed in reimbursement

criteria. Prescriber type categories were: No restriction; Specialist only; None listed; Not available. ‘No restriction’ meant that a healthcare practitioner other than a specialist (e.g. general practitioner) could prescribe interferon-free HCV DAA therapies. ‘Specialist only’ meant that there was a requirement that the prescriber worked within a specialised field. Options included, but were not limited to, hepatologist, infectious disease specialist, internal medicine specialist, and gastroenterologist. If a country permitted a general practitioner to prescribe HCV DAAs but these therapies were not reimbursed, then this was categorised as a restriction (i.e. Specialist only) because we were specifically interested in access to reimbursed therapies. Moreover, the study focus was specific to prescribing; whether a general practitioner or nurse practitioner could treat and monitor a patient who received interferon-free HCV DAA therapy was not recorded. Lastly, HIV co-infection categories were: Prioritised; Eligible; None listed; Not available. ‘Prioritised’ meant that persons with HIV-HCV co-infection were given treatment precedence over persons with HCV mono-infection (i.e. had fewer reimbursement restrictions). ‘Eligible’ meant that persons with HIV co-infection had the same restriction criteria as persons with HCV mono-infection. ‘Other’ was also a possible category for any of the primary outcomes, when appropriate.

Data were organized with descriptive statistics with Microsoft Excel<sup>®</sup> (version 2010, Redmond, USA). With separate Excel spreadsheets, two study authors categorised the outcome criteria of ~15 European countries/jurisdictions each. Once completed, the same two study authors independently cross-checked each other’s categorisation against the documentation. Study authors were then contacted by these two study authors when further clarification was required to resolve any inconsistencies. Following this stage, information was updated as appropriate. Map imagery was created with Tableau Software (version 10.2.2, Seattle, USA).

### *Role of the funding source*

The funding source did not have any input into the study design, data collection, data analysis, interpretation of the data, writing of the report, or the decision to submit the manuscript for publication. The corresponding author had full access to all the data in this study and had final responsibility for the decision to submit the publication.

## FINDINGS

All European countries/jurisdictions (100%, n=35) provided reimbursement for DAAs to treat HCV infection. As of August 1, 2017, the most commonly reimbursed therapy was ombitasvir/paritaprevir/ritonavir  $\pm$  dasabuvir  $\pm$  ribavirin (94%, n=33). Sofosbuvir + daclatasvir  $\pm$  ribavirin (63%, n=22) was the least likely to be reimbursed. Most countries reimbursed sofosbuvir/ledipasvir  $\pm$  ribavirin (89%, n=31) and elbasvir/grazoprevir  $\pm$  ribavirin (91%, n=32), and nearly three-quarters (74%, n=26) reimbursed sofosbuvir/velpatasvir  $\pm$  ribavirin. Some countries, namely Estonia, Latvia, Lithuania, Malta, and Romania, had comparatively fewer therapeutic options than other countries that reimbursed all HCV DAAs [Table 1].

Nearly half of countries/jurisdictions (49%, n=17) required evidence of, at minimum, F2 (METAVIR or equivalent). More specifically, 31% (n=11) of countries/jurisdictions required a minimum stage of F2; 17% (n=6) required  $\geq$ F3; 34% (n=12) had no fibrosis stage restrictions; 14% (n=5) had an additional requirement (e.g. point system that operates regardless of fibrosis stage). One country, Malta, had a minimum fibrosis stage of F4 [Figure 1]. There were circumstances in which fibrosis stage was dependent on genotype. For example, Norway had no fibrosis stage restrictions for genotypes 1 and 4. The clinical recommendation in Norway was also age-dependent. Persons with genotype 3, <40 years, without cirrhosis, were offered peginterferon + ribavirin for 12 weeks as the first line of therapy.<sup>19, 20</sup> In Austria and Switzerland, the minimum fibrosis stage was contingent on the HCV DAA therapy prescribed. For instance, sofosbuvir/ledipasvir  $\pm$  ribavirin and sofosbuvir/velpatasvir  $\pm$  ribavirin had a minimum fibrosis stage of F2 while elbasvir/grazoprevir  $\pm$  ribavirin had no fibrosis stage restrictions.<sup>21-24</sup>

Although 83% (n=29) of countries/jurisdictions had no listed drug or alcohol use restrictions, 17% (n=6) required abstinence of drug and alcohol use prior to treatment (e.g.  $\geq 6$  months abstinence) [Figure 2]. This could include, but was not limited to, toxicological reports (e.g. urine drug screening) every three months to verify abstinence [see appendix p.2]. Hungary, in the case of injection drug use or alcohol use, required a psychiatric consult which supported the capability and compliance for treatment. In Romania, persons with HIV-HCV co-infection had to have a negative drug test to receive reimbursed therapy; this restriction did not seem to apply to persons with HCV mono-infection. In two countries, Liechtenstein and Switzerland, people who inject drugs (PWID) were prioritised for treatment uptake with no fibrosis stage restriction for PWID specifically. In countries in which there were no written guideline regarding substance use, study authors of each respective country emphasised that whether persons were eligible for treatment was ultimately up to physician discretion.

Almost all countries/jurisdictions in Europe (94%; n=33) required a specialist, most often a gastroenterologist, hepatologist, internal medicine specialist, or infectious disease specialist to prescribe HCV DAAs [Figure 3]. England did allow prescribing by a general practitioner although specialist input from a local multi-disciplinary committee was required. Operational Delivery Networks, established by the National Health Service, England, are committees that oversee patient care and for HCV treatment. These specific networks determine who can prescribe HCV DAAs and also, prioritise patients for therapy. In Germany, all general practitioners can prescribe reimbursed HCV DAAs. Certain countries, e.g. France, did allow general practitioners trained in HCV care to monitor patients following initiation of HCV therapy. However, France still required a specialist to prescribe the DAA therapy. Some countries provided a list of specialist prescribers or alternatively, a list of specialist centres – e.g. Austria, Czech Republic, Slovakia, and Slovenia – which patients needed to attend to

receive a specialist prescription. Thus, not all specialists were in a position to prescribe HCV DAAs.

Lastly, the majority of European countries/jurisdictions (97%, n=34) had no additional restrictions for HIV-HCV co-infected persons. Among countries that had minimum fibrosis stage restrictions for HCV monoinfected persons, 26% (n=9) of these countries had no fibrosis stage restrictions for HIV-HCV co-infected persons, and hence, HIV-HCV co-infected persons were prioritised for treatment [Figure 4]. Specifically, in Belgium, Croatia, Czech Republic, Greece, Liechtenstein, Malta, Slovakia, Slovenia, and Switzerland, HIV-HCV co-infected persons were offered HCV treatment regardless of fibrosis stage. In Romania, HIV-HCV co-infected persons needed to provide evidence of a negative drug test.

## DISCUSSION

This study demonstrated some similarities concerning DAA therapy access and reimbursement restrictions in Europe. Restrictions based on specialist prescribing were almost universal with 97% (n=33) of countries requiring a specialist to prescribe DAA therapy. Disease-based restrictions were common with nearly half of countries/jurisdictions (49%, n=17) restricting DAA therapy to persons with significant liver disease ( $\geq$ F2). Further, nearly one-fifth of countries (17%, n=6) required patient abstinence from drug and/or alcohol use to qualify for reimbursed therapies. These restrictions are not in agreement with the 2016 European Association for the Study of Liver (EASL) Recommendations on Treatment of Hepatitis C, which state that all patients without contraindications to therapy should be offered treatment.<sup>12</sup> Additionally, in order to meet WHO targets of eliminating viral hepatitis as a major public health threat by 2030, these restrictions will need to be minimised.<sup>18</sup>

To our knowledge, this is the first study to review HCV DAA reimbursement restrictions in Europe. This study is novel and its primary findings have major potential to influence clinical practice and policy. Similar studies on DAA restrictions have been used in legal and advocacy efforts to remove (or reduce) DAA reimbursement restrictions in Canada and US.<sup>8-10</sup> The follow-up US reimbursement study provided evidence of states lessening (not adding) reimbursement restrictions over a mere two year period.<sup>8, 10</sup> This further highlights the importance of conducting this Europe study to provide a baseline from which to compare DAA access in the upcoming years. A follow-up study on DAA access and restrictions in Europe would be beneficial. There were some limitations in the present study. Given that there was no systematic repository or standardized system for the collation of reimbursement restriction information, we relied on an extensive selection of documents as well as ongoing consultation with study authors to verify data accuracy. This study was limited to written



restrictions and prescribing practices by clinicians might differ from reimbursement criteria. For instance, among countries that have no listed drug use restriction, PWID may be much less likely to be offered DAA therapy. Further, if additional restrictions were reviewed, findings might have revealed even greater reimbursement discrepancies across countries/jurisdictions. In Belgium, for example, an elastography test *and* biomarker score were required to assess fibrosis stage (or alternatively, a liver biopsy).<sup>25</sup> Similarly, countries may also differ on policies concerning retreatment. The present study also did not review intra-country variability. Persons who reside in rural-based areas or do not have private health insurance plans may encounter greater obstacles to reimbursed therapies. Such topics warrant further study. Research on heterogeneity in healthcare systems in Europe, specifically, how medication coverage (with varying deductible and/or co-pay arrangements) impacts treatment access at the patient level is also needed. With respect to reimbursement restrictions, we did not specifically look at the distinction between access and choice. In some countries, the available regimens might have been limited on the basis of restrictions put in place by the payer, which may have limited the selection of DAA therapies that physicians could choose to prescribe. Lastly, timeliness of data was a limitation in this study. While efforts were made to include the most current information, online information and documentation concerning DAA availability and reimbursement restrictions were frequently changing as therapies became approved and governments negotiated new agreements with pharmaceutical companies. Government ministries were not contacted to verify information, which is a noted study limitation. Nevertheless, this study provides an accurate assessment of reimbursement restrictions in Europe as of August 1, 2017.

The considerable heterogeneity in DAA availability and reimbursement restrictions observed in Europe is consistent with previous studies evaluating regional differences in reimbursement

in Canada<sup>9</sup> and the US.<sup>8, 10</sup> Of note, 97% (n=34) of countries/jurisdictions in Europe required a specialist to prescribe DAAs compared to up to 42% of provinces/territories in Canada<sup>9</sup> and 67% of US states.<sup>10</sup> Even in European countries/jurisdictions where primary care providers can prescribe, restrictions are in place through managed care networks (e.g. England) or alternatively, physicians are concerned with medical liability claims (e.g. Germany). In Australia, all medical providers (including primary care and drug/alcohol clinicians) are authorized to prescribe DAA therapy. Less experienced prescribers, however, are required to consult a gastroenterologist/hepatologist or infectious disease specialist in relation to appropriateness of DAA treatment. Under this system, an increasing proportion of DAA prescriptions are completed by primary care and non-specialist providers.<sup>26</sup> Several factors are likely to influence current and future prescriber patterns including involvement of non-specialists. Countries with more concentrated geography (less rural/remote regions), well developed specialist treatment centers with extensive referral pathways, and lower proportions of highly marginalized patients, may not require significant involvement of non-specialists. On the other hand, countries that expand to include non-specialists can be encouraged by evidence which demonstrates that HCV treatment outcomes by primary care providers and specialists are comparable.<sup>27, 28</sup>

The WHO viral hepatitis strategy states that everyone living with viral hepatitis should have access to safe, affordable, and effective care.<sup>18</sup> However, nearly half of European countries (49%, n=17) have restricted DAA reimbursement to persons with advanced liver disease ( $\geq$ F2), which is also inconsistent with the 2016 EASL Recommendations on Treatment of Hepatitis C.<sup>12</sup> Limiting DAA access to patients with more advanced liver disease (and other criteria such as genotype and age as seen in Norway) is a form of prioritisation in the context of high DAA drug pricing related to concerns regarding potential health budget impact, and

presumably, an interim strategy while awaiting development of further treatment infrastructure and declines in DAA pricing. Removal of liver disease stage restrictions is likely to occur in many countries in the near future as has happened recently in France. Successful treatment of HCV infection reduces progression of liver disease<sup>29</sup> and lowers all-cause mortality in people with advanced liver disease.<sup>30</sup> Treatment of those with the greatest risk of transmission (e.g. PWID) also helps to prevent onward HCV transmission.<sup>31</sup> As such, broadened access to DAA therapy will yield both individual and public health benefits.

Seventeen percent (n=6) of European countries restricted access to DAA therapy among people with recent drug and/or alcohol use. Perceptions about poor adherence from ongoing substance use and risk of reinfection due to substance relapse are reasons often put forward to withhold HCV therapy from PWID.<sup>32, 33</sup> A study of HCV practitioners in the DAA era (72% were gastroenterology and hepatology specialists) found that only 15% were willing to treat people who were actively injecting drugs.<sup>32</sup> Promising data have demonstrated excellent adherence and response to DAA HCV therapy among people with recent drug use,<sup>34-39</sup> people receiving opioid substitution therapy,<sup>40-48</sup> and persons with recent injecting drug use.<sup>49-51</sup> Rates of reinfection among PWID are also relatively low.<sup>52, 53</sup> In light of this evidence, there has been some debate as to the ethical justification for withholding therapy from people with ongoing drug/alcohol use.<sup>54</sup>

This study has several key implications for clinicians and policy-makers. European countries should work towards removing restrictions that prevent primary care and drug/alcohol providers from prescribing DAA therapies. The 2016 EASL Recommendations on Treatment of Hepatitis C provide the guidance to ensure appropriate HCV management by medical providers.<sup>12</sup> Mobilising specialists to mentor/train primary care and drug/alcohol providers,

simplifying pathways for the referral of people with advanced liver disease (e.g. APRI >1) to specialists, and providing HCV education and training for nurses, primary care providers, and drug/alcohol providers would all help to enable DAA access. The upcoming availability of pan-genotypic DAA therapies will also facilitate prescribing by primary care and drug/alcohol providers.

While there is no ‘one size fits all’ strategy for the widespread implementation of HCV DAAs, a massive scale-up of testing, linkage to care, and treatment, particularly among high-risk groups, will be required to reduce HCV incidence, HCV prevalence, and HCV-related morbidity and mortality.<sup>5, 17, 55, 56</sup> Future efforts are needed to ensure appropriate monitoring and evaluation of country-level responses to HCV infection (which could then be added to national strategies and action plans) as well as the impact of DAA treatment uptake on HCV-related morbidity and mortality.<sup>18, 57-59</sup> Updated epidemiological data on the morbidity and mortality of HCV-related burden<sup>14, 60</sup> would likely further strengthen national-level policy support for broadened HCV DAA access.

DAA reimbursement restrictions throughout Europe are undoubtedly linked to the list price of DAA regimens. As a result, current reimbursement restrictions exist in most European countries related to prioritisation. The details of DAA prices (or discounts to list prices) are often not readily available, but there are considerable between-country differences in the discounts to list prices across Europe. Broad DAA access requires negotiations to lower DAA prices (or discounts to list prices) to facilitate removal of restrictions. Greater transparency in regard to these negotiations and outcomes is important for the broader strategic development towards “access for all”. The WHO mortality and incidence elimination targets are achievable and cost-effective<sup>61</sup> in many countries but will require the collective efforts of

researchers, healthcare providers, policy-makers, the affected community, and the pharmaceutical industry to succeed.

## **Contributors**

Jason Grebely, Alison D. Marshall, Jeffrey V. Lazarus, Evan B. Cunningham, and Stine Nielsen contributed to study conception and design. A study concept sheet was circulated to all authors who provided comment on the study design. All authors contributed to data acquisition. Alison D. Marshall, Evan B. Cunningham, and Stine Nielsen contacted each study author from the country of interest to facilitate document access. Document searches were also conducted independently by Alison D. Marshall, Evan B. Cunningham, and Stine Nielsen to further corroborate search findings. Alison D. Marshall and Evan B. Cunningham categorized the outcome criteria of ~15 European countries/jurisdictions each and independently cross-checked each other's categorization against the documentation. Alison D. Marshall, Jason Grebely, Gregory J. Dore, Evan B. Cunningham, and Jeffrey V. Lazarus contributed to the drafting of the manuscript. All authors revised the manuscript.

## **Declaration of interests**

Alessio Aghemo has received a research grant from Gilead Sciences, is on the advisory board for Janssen, Merck Sharp & Dohme, Bristol-Myers Squibb, Gilead Sciences, AbbVie, and has received personal fees from Janssen, Merck Sharp & Dohme, Bristol-Myers Squibb, Gilead Sciences, and AbbVie, outside the submitted work. Philip Bruggmann has received grants and personal fees from AbbVie, grants and personal fees from Merck Sharp & Dohme, grants and personal fees from Gilead, and grants and personal fees from Bristol-Myers Squibb, outside the submitted work. Olav Dalgard has received grants from Gilead Sciences, grants and personal fees from AbbVie, grants and personal fees from Merck Sharp & Dohme, outside the submitted work. Carole Seguin-Devaux has received grants from Gilead Sciences, outside the submitted work. Gregory J. Dore has received grants from AbbVie, Merck, Bristol-Myers Squibb, Janssen, and Roche, and personal fees from Gilead Sciences, AbbVie, Merck, Bristol-

Myers Squibb, Janssen, Roche, GlaxoSmithKline, and Abbott Diagnostics, and non-financial support from Gilead Sciences, AbbVie, Merck, Bristol-Myers Squibb, and Roche, outside the submitted work. Robert Flisiak has received grants and personal fees from AbbVie, grants and personal fees from Gilead Sciences, grants and personal fees from Merck, grants and personal fees from Roche, and personal fees from Janssen, outside the submitted work. Graham Foster has received grants and personal fees from Merck, grants and personal fees from Gilead Sciences, and grants and personal fees from AbbVie, during the conduct of the study, and is the National Clinical Lead for Hepatitis C in England. Jason Grebely has received grants from AbbVie, grants from Bristol-Myers Squibb, grants and personal fees from Gilead Sciences, grants and personal fees from Merck, and grants from Cepheid, outside the submitted work. Liana Gheorghe has received personal fees from AbbVie, Bristol-Myers Squibb, Gilead Sciences, and Merck Sharp & Dohme, and is part of the consulting/advisory board for Merck Sharp & Dohme, AbbVie and Gilead Sciences. David Goldberg has received personal fees (honoraria) from Gilead Sciences, AbbVie, and Merck for non-product related lectures, outside the submitted work. Matthew Hickman has received personal fees from Merck Sharp & Dohme, AbbVie, and Gilead Sciences, outside the submitted work. Ligita Jancorienė has received personal fees and non-financial support (consulting fees, honorarium for lectures and payment for conducting clinical studies) from Merck Sharp & Dohme and AbbVie, outside the submitted work. Peter Jarcuska has received personal fees and non-financial support from AbbVie, personal fees and non-financial support from Gilead Sciences, and personal fees from Merck Sharp & Dohme, outside the submitted work. Martin Kåberg has received grants and personal fees from Gilead Sciences, personal fees from AbbVie, and personal fees from Merck Sharp & Dohme, outside the submitted work. Jeffrey V. Lazarus has received research grants and personal fees from AbbVie, Gilead Sciences, and Merck Sharp & Dohme, outside the submitted work. Michael Makara has been an investigator in clinical trials for Novartis,

Bristol-Myers Squibb, Janssen-Cilag, AbbVie, Roche, Boehringer-Ingelheim, Merck Sharp & Dohme, and Regulus, and has received personal fees from Janssen-Cilag, AbbVie, Roche, Boehringer-Ingelheim, Merck Sharp and Dohme, and Gilead Sciences, outside the submitted work. Rui Tato Marinho has received personal fees and advisory board/speaker fees from AbbVie, personal fees and advisory board/speaker fees from Merck Sharp & Dohme, and personal fees and advisory board/speaker fees from Gilead Sciences. Sigurður Ólafsson has received personal fees from Merck Sharp & Dohme, outside the submitted work. Carlos Roncero has received speaker fees from Janssen-Cilag, Ferrer-Brainfarma, Pfizer, Reckitt-Benckiser/Indivior, Lundbeck, Otsuka, Servier, Lilly, GSK, Astra, Sanofi, and Excelsis, received financial compensation for participation as a member of Janseen-Cilag, Indivior, Gilead Sciences, Merck Sharp & Dohme, and Munidipharma Board, carried out grants funded by Reckitt-Benckisert/Indivior, and Gilead Sciences, outside the submitted work. Anne Øvrehus has received personal fees and other (travel, speaker fee and consultancy) from AbbVie, grants, personal fees and other (travel and consultancy) from Gilead Sciences, personal fees from Bristol-Myers Squibb, and other (travel) from Merck Sharp & Dohme, outside the submitted work. James Pocock has received non-financial support from Gilead Sciences and non-financial support from AbbVie, outside the submitted work. Marieta Simonova has received speaker fees from AbbVie, Gilead Sciences, and Merck, and has been an advisor for AbbVie, Gilead Sciences, and Merck, outside the submitted work. Jan Sperl has received grants and personal fees from AbbVie, personal fees from Merck, personal fees from Gilead Sciences, personal fees from Bristol-Myers Squibb, and personal fees from Herbacos Recordati, outside the submitted work. Geert Robaeys has received research grants from Merck Sharp & Dohme, AbbVie, Janssen Pharmaceuticals, and has acted as a consultant/advisor for Gilead Sciences, AbbVie, Merck Sharp & Dohme, and Bristol-Myers Squibb. Ieva Tolmane has received honoraria for lectures from Merck Sharp & Dohme, and



AbbVie. Marc van der Valk has received personal fees from AbbVie, personal fees from Bristol-Myers Squibb, grants and personal fees from Gilead Sciences, personal fees from Johnson & Johnson, grants, personal fees, and non-financial support from Merck Sharp & Dohme, and personal fees from ViiV, outside the submitted work. Adriana Vince has received personal fees and non-financial support from Gilead Sciences, personal fees from Merck Sharp & Dohme, and personal fees from AbbVie during the conduct of this study.

## **Acknowledgements**

Study authors would like to thank the following persons for their assistance with retrieval of documentation and/or interpretation of documentation: Håvard Midgard (Department of Infectious Diseases, Akershus University Hospital, Norway; Institute for Clinical Medicine, University of Oslo, Norway; Department of Gastroenterology, Oslo University Hospital, Norway), Ecaterina Filep (The Kirby Institute, UNSW Sydney, Australia), Gerard Estivill Mercade (The Kirby Institute, UNSW Sydney, Australia), Marcel Schulz (The Kirby Institute, UNSW Sydney, Australia), Rainer Pühr (The Kirby Institute, UNSW Sydney, Australia), Petros Katsioloudes (Ministry of Health, Cyprus), Ioannis Demetriades (Grigorios Clinic, Larnaca General Hospital, Cyprus), the All Wales Therapeutics and Toxicology Centre (Penarth, UK), and the NIHR Health Protection Research Unit (HPRU) in Evaluation of Interventions (University of Bristol, UK).

The Kirby Institute is funded by the Australian Government Department of Health. The views expressed in this publication do not necessarily represent the position of the Australian Government. ADM holds a University International Postgraduate Award from UNSW Sydney and is also supported by the CanHepC Trainee Program (Canada). EBC is supported by the CanHepC Trainee Program (Canada). JG is supported by a National Health and Medical

Research Council Career Development Fellowship. GJD is supported by a National Health and Medical Research Council Practitioner Research Fellowship.

## REFERENCES

1. Polaris Observatory HCV Collaborators. Global prevalence and genotype distribution of hepatitis C virus infection in 2015: a modelling study. *Lancet Gastroenterol Hepatol*. 2017;2(3):161-76.
2. World Health Organization. Global Hepatitis Report 2017. Geneva World Health Organization 2017; License: CC BY-NC-SA 3.0 IG0.
3. Global, regional, and national age–sex specific all-cause and cause-specific mortality for 240 causes of death, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *The Lancet*. 2015;385(9963):117-71.
4. Dore GJ, Feld JJ. Hepatitis C virus therapeutic development: in pursuit of "perfectovir". *Clin Infect Dis*. 2015;60(12):1829-36.
5. Wedemeyer H, Dore GJ, Ward JW. Estimates on HCV disease burden worldwide - filling the gaps. *J Viral Hepat*. 2015;22 Suppl 1:1-5.
6. Gane E, Kershenobich D, Seguin-Devaux C, Kristian P, Aho I, Dalgard O, et al. Strategies to manage hepatitis C virus (HCV) infection disease burden - volume 2. *J Viral Hepat*. 2015;22 Suppl 1:46-73.
7. Nahon P, Bourcier V, Layese R, Audureau E, Cagnot C, Marcellin P, et al. Eradication of Hepatitis C Virus Infection in Patients With Cirrhosis Reduces Risk of Liver and Non-Liver Complications. *Gastroenterology*. 2017;152(1):142-56 e2.
8. Barua S, Greenwald R, Grebely J, Dore GJ, Swan T, Taylor LE. Restrictions for Medicaid Reimbursement of Sofosbuvir for the Treatment of Hepatitis C Virus Infection in the United States. *Ann Intern Med*. 2015;163(3):215-23.
9. Marshall AD, Saeed S, Barrett L, Cooper CL, Treloar C, Bruneau J, et al. Restrictions for reimbursement of direct-acting antiviral treatment for hepatitis C virus infection in Canada: a descriptive study. *CMAJ open*. 2016;4(4):E605-E14.
10. Ooka K, Connolly JJ, Lim JK. Medicaid Reimbursement for Oral Direct Antiviral Agents for the Treatment of Chronic Hepatitis C. *Am J Gastroenterol*. 2017;112(6):828-32.
11. AASLD/IDSA HCV Guidance Panel. Hepatitis C guidance: AASLD-IDSA recommendations for testing, managing, and treating adults infected with hepatitis C virus. *Hepatology*. 2015;62(3):932-54.
12. EASL. EASL Recommendations on treatment of hepatitis C 2016. *J Hepatol*. 2017;66(1):153-94.
13. World Health Organization. Guidelines for the screening, care and treatment of persons with chronic hepatitis C infection 2016. 1-138]. Available from: Available: <http://www.who.int/hepatitis/publications/hepatitis-c-guidelines-2016/en/> (accessed 2017 January 12).
14. Razavi H, Robbins S, Zeuzem S, Negro F, Buti M, Duberg A-S, et al. Hepatitis C virus prevalence and level of intervention required to achieve the WHO targets for elimination in the European Union by 2030: a modelling study. *Lancet Gastroenterol Hepatol*. 2017;2(5):325-36.
15. Hajarizadeh B, Grebely J, Dore GJ. Epidemiology and natural history of HCV infection. *Nat Rev Gastroenterol Hepatol*. 2013;10(9):553-62.
16. Hatzakis A, Chulanov V, Gadano AC, Bergin C, Ben-Ari Z, Mossong J, et al. The present and future disease burden of hepatitis C virus (HCV) infections with today's treatment paradigm - volume 2. *J Viral Hepat*. 2015;22 Suppl 1:26-45.
17. Saraswat V, Norris S, de Knecht RJ, Sanchez Avila JF, Sonderup M, Zuckerman E, et al. Historical epidemiology of hepatitis C virus (HCV) in select countries - volume 2. *J Viral Hepat*. 2015;22 Suppl 1:6-25.

18. World Health Organization. Global health sector strategy on viral hepatitis 2016-2021. Towards ending viral hepatitis. June 2016. 1-56]. Available from: <http://www.who.int/hepatitis/strategy2016-2021/ghss-hep/en/> (accessed 2017 May 21).
19. Sykehusinnkjøp HF. HCV guidelines for the Norwegian Health Authorities. February 8, 2017. . Available from: Available: <http://sykehusinnkjop.no/wp-content/uploads/2017/02/LIS-HCV-anbefalinger-2017-leverand%C3%B8rer.pdf> (accessed 2017 May 10).
20. Norwegian Medical Association. Faglig Veileder for Utredning og Behandling av Hepatitt C hos voksne. March 2017. Clinical Recommendations for Hepatitis C Treatment in Norway. Contact Ministry of Health Norway for more information (accessed 2017 May 5).
21. Schweizerische Eidgenossenschaft. Spezialitätenliste (SL), Index der SL-Verzeichnisse. 2017. Online Drug Formulary. Available from: Available: <http://www.spezialitaetenliste.ch/> (accessed 23 February 2017).
22. Chandrasekhar A. Swiss Hepatitis C Sufferers to get Full Access to Expensive Drug. June 26, 2017. Available from: Available: <https://www.swissinfo.ch/eng/zepatier-swiss-hepatitis-c-sufferers-to-get-full-access-to-expensive-drug/43288924> (accessed 2017 June 27).
23. Erstattungskodex – EKO. Im Blickpunkt-Tipps. NEU im EKO. Sammelband Mai, Juni und Juli 2017. Available from: Available: <http://www.bgkk.at/cdscontent/load?contentid=10008.642874&version=1499412330> (accessed 2017 August 1).
24. Schweizerische Eidgenossenschaft. Drugs Against Hepatitis C: Strong Price Reduction and Extended Remuneration. 2017. Available from: Available: [www.bag.admin.ch/bag/de/home/aktuell/news/news-26-06-2017.html](http://www.bag.admin.ch/bag/de/home/aktuell/news/news-26-06-2017.html) (accessed 2017 June 27).
25. Belgian Association for the Study of the Liver. Belgian HCV Therapy Guidance Update January 2017. Available from: Available: [http://www.basl.be/sites/default/files/Belgian%20HCV%20therapy%20guidance%20update%20january%202017\\_final\\_25012017.pdf](http://www.basl.be/sites/default/files/Belgian%20HCV%20therapy%20guidance%20update%20january%202017_final_25012017.pdf) (accessed 2017 May 19).
26. The Kirby Institute. Monitoring hepatitis C treatment uptake in Australia (Issue 3) Sydney, Australia [updated February 2017]. Available from: Available: [https://kirby.unsw.edu.au/sites/default/files/kirby/report/Monitoring-hep-C-treatment-uptake-in-Australia\\_Iss6-FEB17.pdf](https://kirby.unsw.edu.au/sites/default/files/kirby/report/Monitoring-hep-C-treatment-uptake-in-Australia_Iss6-FEB17.pdf) (accessed 2017 March 17).
27. Arora S, Thornton K, Murata G, Deming P, Kalishman S, Dion D, et al. Outcomes of treatment for hepatitis C virus infection by primary care providers. *N Engl J Med*. 2011;364(23):2199-207.
28. Kattakuzhy SM, Gross C, Teferi G, Jenkins V, Emmanuel B, Masur H, et al., editors. High Efficacy of HCV Treatment by Primary Care Providers: The ASCEND Study. Conference on Retroviruses and Opportunistic Infections; February 22–25, 2016 Boston, MA, USA.
29. Grebely J, Dore GJ. What is killing people with hepatitis C virus infection? *Semin Liver Dis*. 2011;31(4):331-9.
30. van der Meer AJ, Wedemeyer H, Feld JJ, Dufour JF, Zeuzem S, Hansen BE, et al. Life expectancy in patients with chronic HCV infection and cirrhosis compared with a general population. *JAMA*. 2014;312(18):1927-8.
31. Martin NK, Vickerman P, Grebely J, Hellard M, Hutchinson SJ, Lima VD, et al. Hepatitis C virus treatment for prevention among people who inject drugs: Modeling

- treatment scale-up in the age of direct-acting antivirals. *Hepatology*. 2013;58(5):1598-609.
32. Asher AK, Portillo CJ, Cooper BA, Dawson-Rose C, Vlahov D, Page KA. Clinicians' views of hepatitis C virus treatment candidacy with direct-acting antiviral regimens for people who inject drugs. *Subst Use Misuse*. 2016;51(9):1218-23.
  33. Grebely J, Oser M, Taylor LE, Dore GJ. Breaking down the barriers to hepatitis C virus (HCV) treatment among individuals with HCV/HIV coinfection: action required at the system, provider, and patient levels. *J Infect Dis*. 2013;207 Suppl 1:S19-25.
  34. Norton BL, Fleming J, Steinman M, Yu K, Deluca J, Cunningham CO, et al., editors. High HCV Cure Rates for Drug Users Treated with DAAs at an Urban Primary Care Clinic. Conference on Retroviruses and Opportunistic Infections; Feb 22-24 Boston, MA, USA.
  35. Conway B, Raycraft T, Bhutani Y, Kiani G, Shahi R, Singh A, et al. Efficacy of all-oral HCV therapy in people who inject drugs (Abstract #1992). *Hepatology*. 2016;64(S1):990A.
  36. Sulkowski M, Ward K, Falade-Nwulia O, Moon J, Sutcliffe C, Brinkley S, et al. Randomized controlled trial of cash incentives or peer mentors to improve HCV linkage and treatment among HIV/HCV coinfecting persons who inject drugs: the CHAMPS Study. *J Hepatol*. 2017;66:S719.
  37. Mason K, Dodd Z, Guyton M, Tookey P, Lettner B, Matelski J, et al. Understanding Real-World Adherence in the Directly Acting Antiviral Era: a prospective evaluation of adherence amongst people with a history of drug use at a community-based program in Toronto, Canada. *Int J Drug Policy*. 2017;In Press.
  38. Morris L, Smirnov A, Kvassay A, Leslie E, Kavanagh R, Alexander N, et al. Initial outcomes of integrated community-based hepatitis C treatment for people who inject drugs: Findings from the Queensland Injectors' Health Network. *Int J Drug Policy*. 2017.
  39. Read P, Lothian R, Chronister K, Gilliver R, Kearley J, Dore GJ, et al. Delivering direct acting antiviral therapy for hepatitis C to highly marginalised and current drug injecting populations in a targeted primary health care setting. *Int J Drug Policy*. 2017.
  40. Puoti M, Cooper C, Sulkowski MS, Foster GR, Berg T, Villa E, et al. ABT-450/r/Ombitasvir plus Dasabuvir With or Without Ribavirin in HCV Genotype 1-infected Patients Receiving Stable Opioid Substitution Treatment: Pooled Analysis of Efficacy and Safety in Phase 2 and Phase 3 Trials. *Hepatology*. 2014;60:1135a-6a.
  41. Feld JJ, Kowdley KV, Coakley E, Sigal S, Nelson DR, Crawford D, et al. Treatment of HCV with ABT-450/r-ombitasvir and dasabuvir with ribavirin. *N Engl J Med*. 2014;370(17):1594-603.
  42. Zeuzem S, Ghalib R, Reddy KR, Pockros PJ, Ben Ari Z, Zhao Y, et al. Grazoprevir-Elbasvir combination therapy for treatment-naïve cirrhotic and noncirrhotic patients with chronic hepatitis C virus genotype 1, 4, or 6 infection: a randomized trial. *Ann Intern Med*. 2015;163(1):1-13.
  43. Grebely J, Dore GJ, Zeuzem S, Aspinall RJ, Fox R, Han L, et al. Efficacy and safety of sofosbuvir/velpatasvir in patients with chronic hepatitis C virus infection receiving opioid substitution therapy: analysis of phase 3 ASTRAL trials. *Clin Infect Dis*. 2016.
  44. Grebely J, Mauss S, Brown A, Bronowicki JP, Puoti M, Wyles D, et al. Efficacy and Safety of ledipasvir/sofosbuvir with and without ribavirin in patients with chronic HCV genotype 1 infection receiving opioid substitution therapy: analysis of phase 3 ION trials. *Clin Infect Dis*. 2016.

45. Grebely J, Puoti M, Wedemeyer H, Cooper CS, Sulkowski MS, Foster GF, et al. Safety and efficacy of ombitasvir, paritaprevir/ritonavir and dasabuvir with or without ribavirin in chronic hepatitis C patients receiving opioid substitution therapy: a pooled analysis across 12 clinical trials. *J Hepatol.* 2017;66:S514.
46. Grebely J, Jacobson IM, Kayali Z, Verna EC, Shiffman ML, Hyland RH, et al. Sof/vel/vox for 8 or 12 weeks is well tolerated and results in high SVR12 rates in patients receiving opioid substitution therapy. *J Hepatol.* 2017;66:S513.
47. Lalezari J, Sullivan JG, Varunok P, Galen E, Kowdley KV, Rustgi V, et al. Ombitasvir/paritaprevir/r and dasabuvir plus ribavirin in HCV genotype 1-infected patients on methadone or buprenorphine. *J Hepatol.* 2015;63(2):364-9.
48. Dore GJ, Altice F, Litwin AH, Dalgard O, Gane EJ, Shibolet O, et al. Elbasvir-grazoprevir to treat hepatitis C virus infection in persons receiving opioid agonist therapy: a randomized trial. *Ann Intern Med.* 2016;165(9):625-34.
49. Bouscaillou J, Kikvidze T, Butsashvili M, Labartkava K, Inaridze I, Etienne A, et al. Effectiveness of DAA-based treatment of HCV in active people who inject drugs living in middle income countries (MIC): the results of a prospective cohort study in Tbilisi, Georgia. *J Hepatol.* 2017;66:S409.
50. Boglione L, Mornese Pinna S, De Nicolo A, Cusato J, Cariti G, Di Perri G, et al. Treatment with direct-acting antiviral agents of hepatitis C virus infection in injecting drug users: A prospective study. *J Viral Hepat.* 2017.
51. Grebely J, Dalgard O, Conway B, Cunningham E, Bruggmann P, Hajarizadeh B, et al. Efficacy and safety of sofosbuvir/velpatasvir in people with chronic hepatitis C virus infection and recent injecting drug use: the SIMPLIFY study. *J Hepatol.* 2017;66:S513.
52. Dore GJ, Altice F, Litwin AH, Dalgard O, Gane EJ, Shibolet O, et al. Elbasvir/Grazoprevir to Treat HCV Infection in Persons Receiving Opioid Agonist Therapy: A Randomized Controlled Trial (C-EDGE CO-STAR). *Annals of internal medicine.* 2016;In Press.
53. Aspinall EJ, Corson S, Doyle JS, Grebely J, Hutchinson SJ, Dore GJ, et al. Treatment of hepatitis C virus infection among people who are actively injecting drugs: a systematic review and meta-analysis. *Clin Infect Dis.* 2013;57 Suppl 2:S80-9.
54. Grebely J, Haire B, Taylor LE, Macneill P, Litwin AH, Swan T, et al. Excluding people who use drugs or alcohol from access to hepatitis C treatments - Is this fair, given the available data? *J Hepatol.* 2015;63(4):779-82.
55. Liakina V, Hamid S, Tanaka J, Olafsson S, Sharara AI, Alavian SM, et al. Historical epidemiology of hepatitis C virus (HCV) in select countries - volume 3. *J Viral Hepat.* 2015;22 Suppl 4:4-20.
56. Bruggmann P, Berg T, Ovrehus AL, Moreno C, Brandao Mello CE, Roudot-Thoraval F, et al. Historical epidemiology of hepatitis C virus (HCV) in selected countries. *J Viral Hepat.* 2014;21 Suppl 1:5-33.
57. World Health Organization. Manual for the development and assessment of national viral hepatitis plans. September 2015. . 1-62.]. Available from: Available: [www.who.int/hepatitis/publications/manual-hep-plan/en/](http://www.who.int/hepatitis/publications/manual-hep-plan/en/) (accessed 2017 May 19)
58. European Liver Patients Association. The 2016 Hep-Core Report. Monitoring the implementation of hepatitis B and C policy recommendations in Europe. March 2017. 1-158]. Available from: [http://www.elpa.eu/sites/default/files/project-documents/Hep-CORE\\_full\\_report\\_21Mar2017\\_errata\\_Final.pdf](http://www.elpa.eu/sites/default/files/project-documents/Hep-CORE_full_report_21Mar2017_errata_Final.pdf) (accessed 2017 May 30).
59. World Health Organization. Regional committee for Europe 66th session. Action plan for the health sector response to viral hepatitis in the WHO European Region. Copenhagen, Denmark, 12-15 September 2016, . Available from: Available:

<http://www.euro.who.int/en/about-us/governance/regional-committee-for-europe/66th-session/documentation/working-documents/eurrc6610-action-plan-for-the-health-sector-response-to-viral-hepatitis-in-the-who-european-region> (accessed 2017 June 23).

60. Muhlberger N, Schwarzer R, Lettmeier B, Sroczynski G, Zeuzem S, Siebert U. HCV-related burden of disease in Europe: a systematic assessment of incidence, prevalence, morbidity, and mortality. *BMC Public Health*. 2009;9:34.
61. Scott N, McBryde ES, Thompson A, Doyle JS, Hellard ME. Treatment scale-up to achieve global HCV incidence and mortality elimination targets: a cost-effectiveness model. *Gut*. 2016.





**Table 1. Availability of reimbursed interferon-free DAAs for HCV infection in Europe**

	HCV DAA Therapy					
	SOF + RBV	SOF/LED ± RBV	SOF/VEL ± RBV	P-OD ± RBV	EBR-GZR ± RBV	SOF + DCV ± RBV
Austria	YES	YES	YES	YES	YES <sub>μ</sub>	YES
Belgium	YES	YES	YES	YES	YES	YES
Bulgaria	YES	YES	NO	YES	YES	NO
Croatia	YES	YES	YES	YES	YES	NO
Cyprus	YES	YES	YES	YES	YES	NO
Czech Republic	YES	YES	YES	YES	YES	YES
Denmark	YES	YES	YES	YES	YES	YES
England	YES	YES	YES	YES	YES	YES <sup>β</sup>
Estonia	NO	NO	NO	YES <sup>‡</sup>	YES <sup>‡</sup>	NO
Finland	YES	YES	YES	NO	YES	NO
France	YES	YES	YES	YES	YES	YES
Germany	YES	YES	YES	YES	YES	YES
Greece	YES	YES	YES	YES	YES	YES
Hungary	NO	YES	NO	YES	NO	NO
Iceland	YES	YES	YES	YES	YES	YES
Ireland	NO	YES	YES	YES	YES	YES
Italy	YES	YES	YES	YES	YES	YES
Latvia	NO	NO	NO	YES <sup>β</sup>	YES <sup>β</sup>	NO
Liechtenstein	YES	YES	YES	YES	YES <sup>μ</sup>	YES
Lithuania	NO	NO	NO	YES	YES <sup>β</sup>	NO
Luxembourg	YES	YES	YES	YES	YES	YES
Malta	YES	YES	NO	NO	NO	NO
Netherlands	YES	YES	YES	YES	YES	YES
N. Ireland	YES	YES	YES	YES	YES	YES
Norway	YES	YES	YES	YES	YES	YES
Poland	YES	YES	NO	YES	YES	NO
Portugal	YES	YES	YES	YES	YES	YES
Romania	NO	NO	NO	YES	NO	NO
Scotland	YES	YES	YES	YES	YES	YES <sup>†</sup>
Slovakia	YES	YES	NO	YES	YES	NO
Slovenia	YES	YES	YES	YES	YES	NO
Spain	YES	YES	YES	YES	YES	YES
Sweden	YES*	YES*	YES	YES	YES	YES*
Switzerland	YES	YES	YES	YES <sub>μ</sub>	YES <sub>μ</sub>	YES
Wales	YES	YES	YES	YES	YES	YES <sup>†</sup>

<sup>β</sup>Restricted to fibrosis stage of ≥F3

\*Special consideration is required

<sup>‡</sup>Need to be GT3, treatment naïve with ≥F3

<sub>μ</sub>There are no fibrosis stage restrictions for GT1 and GT4

<sup>‡</sup>Reimbursed regardless of fibrosis stage in patients with cryoglobulinemia and post orthotopic transplantation

- <sup>a</sup> Fibrosis stage restrictions based on HCV genotype
- <sup>b</sup> Fibrosis stage is included in a point system for prioritisation of DAA therapy
- <sup>c</sup> Fibrosis stage restrictions based on HCV genotype and IL28B polymorphism
- <sup>d</sup> Fibrosis stage restrictions based on

**Figure 1. Minimum fibrosis stage required for reimbursement of interferon-free DAAs for treatment naïve patients with HCV infection in Europe**

\*Please note: This figure was uploaded separately

<sup>a</sup>  
Persons with HIV-HCV co-infection had to have a negative drug test in order to receive reimbursed therapy

**Figure 2. Drug and alcohol restrictions for reimbursement of interferon-free DAAs for patients with HCV infection in Europe**

\*Please note: This figure was uploaded separately

**Figure 3. Prescriber-type restrictions for reimbursement of interferon-free DAAs for patients with HCV infection in Europe**


\*Please note: This figure was uploaded separately

<sup>a</sup> Persons with HIV-HCV co-infection had to have a negative drug test in order to receive reimbursed therapy


**Figure 4. HIV co-infection restrictions for reimbursement of interferon-free DAAs for patients with HCV infection in Europe**

\*Please note: This figure was uploaded separately


Figure



Figure




Figure





Figure



**Necessary Additional Data**

[Click here to download Necessary Additional Data: Supplementary Appendix\\_August 18 2017.pdf](#)

[Click here to download Necessary Additional Data: drgalc wo discretion.emf](#)

drgalc.pdf

[Click here to download Necessary Additional Data: Drgalc.pdf](#)

[Click here to download Necessary Additional Data: Fibrosis stage dash.emf](#)

**fibrosis.pdf**

**[Click here to download Necessary Additional Data: Fibrosis.pdf](#)**

[Click here to download Necessary Additional Data: HIV dash.emf](#)

hiv.pdf

[Click here to download Necessary Additional Data: HIV.pdf](#)



[Click here to download Necessary Additional Data: Specialist dash.emf](#)

specialist.pdf

[Click here to download Necessary Additional Data: Specialist.pdf](#)